

# Maintaining internal representations: the role of the human superior parietal lobe

Daniel M. Wolpert<sup>1</sup>, Susan J. Goodbody<sup>1</sup> and Masud Husain<sup>2</sup>

<sup>1</sup> Sobell Department of Neurophysiology, Institute of Neurology, Queen Square, University College London, London WC1N 3BG, UK

<sup>2</sup> Division of Clinical Neuroscience and Psychological Medicine, Imperial College School of Medicine, Charing Cross Hospital, London W6 8RF, UK

Correspondence should be addressed to D.M.W. (wolpert@hera.ucl.ac.uk)

In sensorimotor integration, sensory input and motor output signals are combined to provide an internal estimate of the state of both the world and one's own body. Although a single perceptual and motor snapshot can provide information about the current state, computational models show that the state can be optimally estimated by a recursive process in which an internal estimate is maintained and updated by the current sensory and motor signals. These models predict that an internal state estimate is maintained or stored in the brain. Here we report a patient with a lesion of the superior parietal lobe who shows both sensory and motor deficits consistent with an inability to maintain such an internal representation between updates. Our findings suggest that the superior parietal lobe is critical for sensorimotor integration, by maintaining an internal representation of the body's state.

Damage to the human posterior parietal cortex (PPC) can cause perceptual and motor deficits, including visual mislocalization, neglect or unawareness of contralesional visual or tactile stimuli, impaired perception of contralesional body parts and misreaching to visual objects<sup>1-4</sup>. The study of these syndromes, together with more recent functional imaging investigations of individuals without brain injury<sup>5-8</sup> has led to proposals that the human PPC is important for spatial perception, association of sensory signals, directing attention, visuomotor control and motor planning<sup>4,9-12</sup>. Unfortunately, no unifying theoretical framework has emerged from human studies for understanding the function of the human PPC. Instead, there has been considerable reliance on neurophysiological investigations in monkeys<sup>9,13-17</sup>, and the models developed from them, to formulate working hypotheses of human PPC function. Such extrapolations are, at best, indirect.


We propose a new theoretical framework for the visuomotor functions of the human parietal cortex. Recent developments in the field of sensorimotor integration have provided a new approach to understanding how sensory and motor signals are combined to provide an estimate of the state of both the world and one's own body<sup>18,19</sup>. The approach we use comes from the powerful observer framework in engineering<sup>20</sup>. This framework is relevant when there is a system under control, in our case the body, and the goal is to estimate the state of the system. By state we refer to a compact set of variables that captures both the configuration of the body, such as joint angles or hand position, and its interactions with the world, such as contact with an object. For optimal (most accurate) state estimates, the observer framework requires a system that monitors both the inputs and outputs of the system, in our case the sensory feedback and motor commands. The observer then uses these values to produce online state estimates, which are updated as further

sensory and motor signals arrive. The major objective of the observer is to reduce the uncertainty in the state estimate that arises from noise inherent in both the sensory and motor signals. Integrating these signals over time, rather than relying on a single perceptual or motor snapshot, produces an improved estimate. The new estimate of state is made by recursively updating the current state estimate based on the incoming motor and sensory signals.

A key feature of the observer model is that it requires that the state representation be stored and updated as new information arrives (Fig. 1). Updating the state estimate and storing the state estimate between updates can be considered as two functionally and possibly anatomically distinct processes. Although such an internal state estimate has been proposed within the central nervous system<sup>19,21</sup>, there have been no reports of lesions that result in a specific inability to maintain such an estimate. The key claim in this paper is that the state estimate is stored in the parietal cortex.

## Results

We have studied a patient with a parietal lesion who has unusual sensory and motor deficits that become apparent only as the duration of a sensory stimulus or movement increases. The patient, PJ, is a right-handed, fifty-year-old woman who was first seen two years before our study by the Neurology service following episodes of involuntary jerking of the right arm, which were considered to be focal seizures. Magnetic resonance imaging at that time demonstrated an extra-axial cyst encroaching upon the cortex and subcortical white matter of the left superior parietal lobe. The images are consistent with those of an arachnoid cyst, but similar appearances may occur following trauma. The patient recalls that five years before these fits she sustained an accidental injury to the left side of her head with loss of con-



**Fig. 1.** A schematic of the sensorimotor integration process. The 'observer' monitors and integrates both the inputs and outputs of the system, the sensory feedback and motor commands, by recursively updating the stored state estimate.

sciousness for thirty minutes. At the time she presented with focal seizures affecting the right arm, she declined cystoperitoneal shunting but agreed to regular follow-up scanning. She started on carbamazepine for prevention of further focal seizures. On a maintenance dose of 600 mg per day, she had no recurrence.


Two years later, she has returned complaining of new symptoms. She now reports that she perceives her right arm and leg to drift and then fade unless she is able to see them. For example, while lying in bed, she might realize that she has 'lost' her right arm. Only when she looks at her arm does she know where it is. Similarly, she can be sitting on a bus and find that another passenger has tripped over her right foot, which is situated, without her knowledge, in the middle of the aisle. Repeat magnetic resonance imaging now demonstrates that the cyst has grown radially by 2.5 mm (Fig. 2). Unlike other parietal cases, in which patients are permanently unaware of their limbs, her symptoms are novel because she only becomes unaware of her right arm and leg over tens of seconds. Furthermore, she can regain awareness of the location of her right arm and leg by looking at them, although she cannot voluntary regain awareness otherwise.

On examination, her visual fields were full to confrontation (normal). There was no visual extinction or neglect as assessed by line bisection, cancellation on the Mesulam shape cancellation task<sup>22</sup> and object drawing. Saccadic and smooth-pursuit eye movements seemed normal. Tone and power in the upper and lower limbs were normal bilaterally. Deep tendon reflexes were slightly brisker in the right limbs than the left. Plantar responses were flexor (normal). Although there was subjective diminution of light touch in the right arm, her ability to discriminate sharp from blunt tactile stimuli was good. However, if a tactile stimulus was held still on her right hand, when her eyes were closed, she reported that the sensation faded over seconds until she could no longer detect the stimulus. Proprioception and vibration were also subjectively considered to be different in the right upper limb than the left, but there seemed to be no objective difference on clinical testing. She had right tactile extinction. Astereognosis (impaired ability to recognize objects by their size, shape, and texture) and agraphesthesia (inability to identify numbers written on the palm) were detected in the right hand. When asked to point to

the examiner's finger with free vision, she had no difficulty using either left or right hand. However, when she was asked to fixate centrally on the examiner's nose and point slowly to his finger, she consistently missed when using her right hand, regardless of whether reaches were made into the left or right visual field. There was no limb dyspraxia (motor deficits which are not a consequence of weakness or sensory loss), and her gait was normal.

We investigated the time course of PJ's tactile perception by placing weights on the back of her hand. Although PJ was able to detect, without vision, when an object was placed on the back of her right hand, the percept then faded until she could no longer detect the object (Methods). The time to fade increased linearly ( $r^2 = 0.63$ ,  $p < 0.001$ ) with the mass of the object (Fig. 3). For the lightest weight of 10 grams, the percept took 2.9 seconds to disappear, whereas for a weight of 150 grams, the time to fade increased to 10.5 seconds. Although PJ could no longer detect the object at the end of each trial, she could always detect its removal. If the weight was moved to another part of her hand, she could both detect its removal and its replacement. However, once the weight was stationary, the percept of the weight would eventually disappear. In contrast, when the weight was applied to her left hand, the percept did not fade, and she never reported being unable to detect the object. When vibratory stimuli were applied, the percept did not fade on either hand. Whereas parietal patients with central deafferentation are unable to perceive either constant or changing stimuli<sup>23,24</sup>, PJ becomes effectively deafferented only after a period of several seconds in the presence of a constant tactile stimulus.

To assess the motor consequences of her tactile fading, we examined PJ's ability to maintain a precision grip with and without visual feedback of her performance (Methods). She held a force transducer with her thumb and index finger in a precision grip and was asked to maintain a constant grip force. Grip force was displayed on an oscilloscope screen together with the target level of grip force. Having matched the grip force under visual feedback of the oscilloscope screen, PJ was asked to maintain the grip force constant with either full vision or no vision. When given visual feedback of her performance, PJ was able to maintain her grip level using either hand (Fig. 4a and b). Like neurologically normal subjects<sup>24</sup>, without visual feedback, PJ was still able to maintain her force level using her unaffected left hand (Fig. 4c). However, when using her right hand, the grip force decayed to near zero over 10–15 seconds (Fig. 4d; the removal of



**Fig. 2.** The lesion found in the patient P. T2-weighted axial (left) and T1-weighted coronal (right) magnetic resonance imaging sections demonstrating a left superior parietal cyst.





weight was placed on the back of her hand. The base of the weight was covered by a plastic disc to prevent temperature cues. PJ was asked to report when she could no longer feel the weight on the back of her hand, and the time was recorded. Five different weights (10, 20, 50, 100 and 150 g) were placed on her hand, each for 9 repetitions in a pseudorandom order. A small vibrator was used to examine whether dynamic stimuli faded.

To examine the consequences of the sensory fading, we determined PJ's ability to maintain a constant force output. PJ held a force transducer (Assurance Technologies) with her thumb and index finger 1 cm apart. The grip force produced was sampled at 250 Hz by a CED 1401plus (Cambridge Electronic Design). Grip force was recorded and simultaneously displayed on an oscilloscope screen. Also shown on the oscilloscope was a target level of grip force (either 4.5 or 9 N), which PJ was required to match. Having matched the grip force under visual feedback of the oscilloscope screen, PJ was required to maintain the grip force constant for a further 20 seconds under two different conditions. In the full vision condition, PJ had vision of both her own hand and the oscilloscope displaying the actual and target grip force. In the second, no vision condition, the Plato glasses prevented all visual feedback. Each of these four conditions (two force levels and two visual feedback conditions) was repeated twice for both the right and left hands. Results for the 4.5 N force level (not shown) were similar to those for the 9 N condition.

To quantify the perceived location of her right limb in the absence of visual feedback, PJ was required to track the perceived position of her right hand with her left hand. PJ was seated at a table with her hands placed palms down next to each other, with index fingers parallel. The positions of the fingers were recorded using an active infra-red tracking system (Optotrak 3020, Northern Digital) sampling at 250 Hz. At the beginning of each trial, PJ was instructed to look at the positions of her hands. Vision was then occluded, and PJ tracked the perceived position of the right hand with her left hand. The right hand remained stationary throughout the trial. Three trials for each of four different starting locations of her right hand were recorded for up to 80 seconds.

To quantify PJ's optic ataxia, we assessed her pointing accuracy to visual targets in her periphery by asking her to point, as accurately as possible, to the experimenter's finger while fixating on his nose. Movements were made with both the right and left hand at a self-paced and slow speed. The locations of both the experimenter's and PJ's fingers were recorded using the infrared tracking system. Ten repetitions to different targets were made with both hands and at both speeds.

## Acknowledgements

We thank PJ for participating in the study and Jon Driver and Tobe Freeman for helpful discussions. This work was supported by a grant from the Wellcome Trust and the Royal Society.

RECEIVED 20 MAY; ACCEPTED 22 AUGUST 1998

- Bálint, R. Seelenlähmung des 'schauens', optische ataxie, räumliche störung der aufmerksam. *Monatsh. Psychiat. Neurol.* 25, 51–81 (1909).
- Holmes, G. Disorders of visual orientation. *Br. J. Ophthalmol.* 2, 449–468 (1918).
- McFie, J., Piercy, F. J. & Zangwill, O. L. Visual-spatial agnosia associated with lesions of the right cerebral hemisphere. *Brain* 73, 167–170 (1950).
- DeRenzi, E. *Disorders of Space Exploration and Cognition* (John Wiley, Chichester, 1982).
- Corbetta, M., Miezin, F. M., Shulman, G. L. & Petersen, S. E. A PET study of visuospatial attention. *J. Neurosci.* 13, 1202–1226 (1993).
- Corbetta, M., Shulman, G. L., Miezin, F. M. & Petersen, S. E. Superior parietal cortex activation during spatial attention shifts and visual feature conjunction. *Science* 270, 802–805 (1995).
- Grafton, S. T., Fagg, A. H., Woods, R. P. & Arbib, M. A. Functional-anatomy of pointing and grasping in humans. *Cereb. Cortex* 6, 226–237 (1996).
- Clower, D. M. *et al.* Role of posterior parietal cortex in the recalibration of visually guided reaching. *Nature* 383, 618–621 (1996).
- Andersen, R. in *Handbook of Physiology* vol. V Part II (eds Plum, F. & Mountcastle, V. B.) 483–518 (American Physiological Society, Rockville, Maryland, 1987).
- Milner, A. D. & Goodale, M. *The Visual Brain in Action* (Oxford Univ. Press, Oxford, 1995).
- Jeannerod, M. *The Cognitive Neuroscience of Action* (Blackwell, Oxford, 1997).
- Husain, M. in *Vision and Visual Dysfunction* vol. 13 (ed. Stein, J. F.) 12–43 (Macmillan, Basingstoke, UK, 1991).
- Mountcastle, V. B., Lynch, J. C., Georgopoulos, A. P., Sakata, H. & Acuna, C. Posterior parietal association cortex of the monkey: command function for operations within extrapersonal space. *J. Neurophysiol.* 38, 871–907 (1975).
- Bushnell, M. C., Goldberg, M. E. & Robinson, D. L. Behavioural enhancement of visual responses in monkey cerebral cortex. I. modulation in posterior parietal cortex related to selective visual attention. *J. Neurophysiol.* 46, 755–772 (1981).
- Duhamel, J. R., Colby, C. L. & Goldberg, M. E. The updating of the representation of visual space in parietal cortex by intended eye movements. *Science* 255, 90–92 (1992).
- Andersen, R. A., Snyder, L. H., Bradley, D. C. & Xing, J. Multimodal representation of space in the posterior parietal cortex and its use in planning movements. *Annu. Rev. Neurosci.* 20, 303–330 (1997).
- Rizzolatti, G., Fogassi, L. & Gallese, V. Parietal cortex: from sight to action. *Curr. Opin. Neurobiol.* 7, 562–567 (1997).
- Wolpert, D. M., Ghahramani, Z. & Jordan, M. I. Are arm trajectories planned in kinematic or dynamic coordinates? An adaptation study. *Exp. Brain Res.* 103, 460–470 (1995).
- Wolpert, D. M. Computational approaches to motor control. *Trends Cogn. Sci.* 1, 209–216 (1997).
- Goodwin, G. C. & Sin, K. S. *Adaptive Filtering Prediction and Control* (Prentice-Hall, Englewood Cliffs, New Jersey, 1984).
- Wolpert, D. M., Ghahramani, Z. & Jordan, M. I. An internal model for sensorimotor integration. *Science* 269, 1880–1882 (1995).
- Mesulam, M. M. *Principles of Behavioural Neurology: Tests of Directed Attention and Memory* (Davis FA, Philadelphia, 1985).
- Jeannerod, M., Michel, F. & Prablanc, C. The control of hand movements in a case of hemianaesthesia following a parietal lesion. *Brain* 107, 899–920 (1984).
- Rothwell, J. C. *et al.* Manual motor performance in a deafferented man. *Brain* 105, 515–542 (1982).
- Wann, J. P. & Ibrahim, S. F. Does limb proprioception drift. *Exp. Brain Res.* 91, 162–166 (1992).
- Perenin, M. T. & Vighetto, A. Optic ataxia: a specific disruption in visuomotor mechanisms. I. different aspects of the deficit in reaching for objects. *Brain* 111, 643–674 (1988).
- Critchley, M. *The Parietal Lobes* (Edward Arnold, London, 1953).
- Mennemeier, M. S. *et al.* Contributions of the parietal and frontal lobes to sustained attention and habituation. *Neuropsychologia* 32, 703–716 (1994).
- Holliday, I. E., Kennard, C. & Ruddock, K. H. Rapid fading of visual sensations in a subject with a parietal occipital tumor. *Ophthalmic Physiol. Opt.* 5, 149–156 (1985).
- Head, H. & Holmes, G. Sensory disturbances from cerebral lesions. *Brain* 34, 102–213 (1912).
- Vallar, G. & Perani, D. The anatomy of unilateral neglect after right-hemisphere stroke lesions. A clinical/CT-scan correlation study in man. *Neuropsychologia* 24, 609–622 (1986).
- Mattingley, J. B., Husain, M., Rorden, C., Kennard, C. & Driver, J. Motor role of human inferior parietal lobe revealed in unilateral neglect patients. *Nature* 392, 179–182 (1998).
- Posner, M. I., Walker, J. A., Friedrich, F. J. & Rafal, R. D. Effects of parietal injury on covert orienting of attention. *J. Neurosci.* 4, 1863–1874 (1984).
- Pardo, J. V., Fox, P. T. & Raichle, M. E. Localization of a human system for sustained attention by positron emission tomography. *Nature* 349, 61–64 (1991).
- Coull, J. T., Frith, C. D., Frackowiak, R. S. J. & Grasby, P. M. A frontoparietal network for rapid visual information-processing: a PET study of sustained attention and working-memory. *Neuropsychologia* 34, 1085–1095 (1996).
- Friedrich, F. J., Egly, R., Rafal, R. D. & Beck, D. Spatial attention deficits in humans: a comparison of superior parietal and temporal-parietal junction lesions. *Neuropsychologia* 12, 193–207 (1998).
- Milner, A. D. in *Parietal Lobe Contributions to Orientation in 3D Space* (eds Thier, P. & Karnath, H. O.) 3–22 (Springer-Verlag, Heidelberg, 1997).